

Short note

Bryozoan encrusted plastic from the continental slope: eastern South Island, New Zealand

Rowan Carter¹ and Murray R. Gregory^{1,2}

¹Department of Geology, The University of Auckland, Private Bag 92019
Auckland, New Zealand

²Corresponding author; e-mail: m.gregory@auckland.ac.nz

(Received 8 August 2005, revised and accepted 26 September 2005)

Abstract

A small flake of plastic sheeting recovered from the upper continental slope to the east of the South Island, and upon which was growing a colony of the celeporid bryozoan *Galeopsis mimicus*, is symptomatic of broader environmental problems associated with marine debris accumulation on the deep sea floor.

Keywords: encrusting bryozoan - marine debris - plastic litter - deep sea floor - New Zealand.

Introduction

Over the past 30 years there has been much documentation of the extent to which oceanic surface waters have become increasingly contaminated by plastics and other persistent synthetic materials (i.e., marine debris). Research has largely focussed on quantifying amounts and distribution patterns, establishing types of litter and their sources, evaluation of environmental, aesthetic and commercial impacts, and finding (or speculating about) solutions. The problems are particularly manifest in coastal waters and along shorelines. These aspects, together with possible answers, are comprehensively reviewed in Coe and Rogers (1997). The commonly identified

sinks for this plastic litter are either burial in beach and backshore settings by drifting sand, or fragmentation through photodegradation-induced embrittlement into fine powders that pass from view and are assimilated into the environment. It is also accepted that onshore winds may disperse plastic sheeting and other light material, including foamed polyethylene and polystyrene, across coastal hinterlands.

Although Williams *et al.* (1993) and Goldberg (1997) have recently emphasized that the sea floor, from intertidal and shallow sub-littoral zones to greater depths, is an important sink for marine debris, aspects of its significance in all these settings have been appreciated for some time (e.g., Holmström 1975, Harms 1990). Over

the past few years, records of plastics and other anthropogenic debris accumulating on the sea floor at varying depths have increased. The localities include Antarctica (Lenihan *et al.* 1990); the Bay of Biscay and elsewhere in European waters (Galgani *et al.* 1995a, Galgani *et al.* 2000); the western (Galgani *et al.* 1995b, Galgani *et al.* 1996) and eastern Mediterranean (Bingel *et al.* 1987, Galil *et al.* 1995, Stefatos *et al.* 1999); Alaska (Hess *et al.* 1999); California (Moore & Allen 2000); Indonesia (Uneputty & Evans 1997); Japan (Kanehiro *et al.* 1996) and South Africa and New Zealand (Gregory & Ryan 1997, Backhurst & Cole 2000). The above studies typically record larger (macro- megascopic), readily visible plastic items. Thompson *et al.* (2004), have recently demonstrated that quantities of microscopic plastic fragments and fibres associated with plankton in surface waters have significantly increased since the 1960's, and also that these materials are dispersed through the water column. Such widely distributed occurrences suggest environmental impacts whose magnitude and significance have yet to be fully appreciated (see Goldberg 1997) or explored (e.g., Thompson *et al.* (2004).

While diverse biotas of fouling organisms and encrusters are known from pelagic (i.e., floating) plastics (e.g., Winston 1982, Klauswitz 1984, Ye and Andrady 1991, Stevens *et al.* 1996, Winston *et al.* 1997), similar studies of benthic (i.e., that which has settled on to the sea floor) materials are limited (e.g., Holmström 1975, Harms 1990, Powlik 1995). The scant evidence presently available, however, suggests typical temperate hard ground communities characterised by bryozoans, hydrozoans, sponges and foraminifera, as well as barnacles, bivalves and polychaetes, form

the basis for epibiontic communities. At shallow depths in the photic zone, crustose (coralline) red algae together with soft brown and green algae are often common. Bryozoans are generally a conspicuous, if not dominant, epibiont of both pelagic and benthic marine debris (e.g., Stevens 1992, Barnes & Sanderson 2000).

Material

While picking benthic and planktic foraminifera in bottom sediment samples taken from a number of New Zealand Oceanographic Institute [NZOI now NIWA] grab and core stations, as well as at several ODP sites, lying in water depths between 90 m and 4600 m to the east of Banks Peninsula, South Island, New Zealand and across the Chatham Rise (Figure 1), small flakes of frayed and shredded plastic sheeting were encountered on two occasions. One of these came from the surficial 1-2 cm of




Figure 1. Central New Zealand, showing local sampling stations referred to in the text. (inset: northern Norfolk Ridge and South Loyalty Basin stations, DW33, 36, 37 and 76; CP67 and 75; see Gordon and d'Hondt, 1989).




Figure 2. The cheilostome bryozoan *Galeopsis mimicus* on the frayed plastic substrate (arrowed) to which it is attached (A) and enlarged showing characteristic surface evident (arrowed) (B). Scale bar 200 µm.

an ODP Leg 181 core taken on the upper continental slope at site 1119 (position $44^{\circ} 45.33'S$ / $172^{\circ} 23.60'E$), located in the South Canterbury Bight, some 60 km offshore and at a water depth of 393 m. The flake's dimensions are c.9 mm x 6 mm. This flake is worthy of note because of a small celoporid bryozoan that was attached to it (Figure 2). Despite being somewhat abraded the specimen has been tentatively identified as *Galeopsis mimicus* Gordon 1989. Previous records of this taxon in New Zealand waters are at depths of between 297 m and 520 m a short distance off the coast of Westland (NZOI stations Q693 and 694, S374; Figure 1) where it has been found only on the aboral spines of the frontal notch of the echinoid *Spatangus multispinus* Mortensen (Gordon 1989). It is also known in waters >2000 km to the north of New Zealand, where it occurs between depths of 470 m and 825 m from six stations immediately south of New Caledonia and lying between latitudes $24^{\circ} 55'S$ and $22^{\circ} 18'S$ along the South Loyalty Basin and northern Norfolk Ridge. Here it occurs on axes of small gorgonians (Gordon & d'Hondt 1997). This species seems to like thin structures to settle on and a plastic

flake could be ideal (Gordon, pers comm. 2000). The recorded depth range for *G. mimicus* suggests that it colonised the plastic substrate while it lay on the sea floor and that it is unlikely to have been transported from shallower, up-slope depths. It can also be noted that this is the first record of *G. mimicus* in waters to the east of New Zealand. Dredgings from this region have not, however, been extensively sampled for bryozoans.

In 66 samples examined, the only other plastic artifacts to be found came from NZOI station Q859 (water depth 3654 m: position $39^{\circ} 56.24'S$ / $178^{\circ} 30.48'E$) (Figure 1). These were thread-like scrapings, devoid of any bryozoan encrustation or evidence of attachment scarring, and careful examination suggests contamination from the plastic liner during the core-splitting process.

Discussion

This fortuitous discovery is a further example of plastic reaching the deep sea floor and, once there, providing a suitably substantial or firm substrate for colonisation by fouling epibionts. In general, sheet and other plastic products

are positively to neutrally buoyant, and that significant quantities can reach the deep sea floor is something that many find surprising (e.g., Holmström 1975). The mechanisms by which they do so are poorly understood. For instance, an example is Oshima's (2000) observation of a fleet of flimsy white plastic, supermarket shopping bags, up-ended and suspended at depths of 2000 metres, and drifting like an assembly of ghosts. From experimental evidence, Ye and Andrady (1991) and Stevens (1992) have argued that rapid and heavy fouling of plastic materials may lead to density increases sufficient to permanently sink them. These authors also recognised that cleaning of fouled surfaces by grazers can lead to cyclic episodes of submergence and resurfacing until permanent settlement to the seafloor takes place. It is also possible that plastic sheeting and other items may attract or act as a passive collection surface for non-living particulate detritus. This, together with photodegradation and embrittlement leads to the density increases necessary to take plastic items to the sea-bed (Powlik 1995) without invoking down-welling and entrainment processes.

The presence of plastic sheeting and other large plastic items, as well as discarded fishing gear on the sea floor is undesirable and likely to be environmentally damaging (Williams *et al.* 1993, Goldberg 1997). It must however be recognised that hard grounds, even small-scale ones, will attract a biota different from that of adjacent soft, muddy substrates. It seems ironic to suggest that plastic artifacts reaching the seafloor could enhance local biological richness and diversity in the short term, although in the long term this material is doomed for permanent entombment in

slowly accumulating sediment.

Conclusions

The occurrence of *Galeopsis mimicus* on a flake of plastic sheeting recovered from the deep seafloor may be viewed by some as little more than an interesting curiosity, similar to the beach-cast virgin plastic granule from northern New Zealand that hosted the bryozoan *Membranipora tuberculata* which was illustrated by figure 8 in Gregory (1978). It was inferred that arrival of the latter taxon was by way of eastwards dispersal from Australian waters and across the northern Tasman Sea in eddies of the East Australian Current (Gregory 1978). Both, however, can be considered to be evidence symptomatic of much wider problems, and ones having potentially serious environmental implications. Passively drifting, discarded plastic materials may be colonized by a diverse community of encrusting and fouling epibionts (Stevens 1992, Winstone *et al.* 1997). This cargo of hitch-hikers and hangers-on may include aggressive and invasive marine species. Rafting of alien taxa rafted in this way could endanger environments both near and far from their natural habitats (Winston 1997, Gregory 1998, 2004). Similarly, plastic debris could be a vector effecting the introduction of aggressive alien taxa into regions that hitherto have been considered impenetrably deep.

Acknowledgements

This note was completed using facilities in the Geology Department and the Research Centre for Surface and Materials Science at the University of Auckland. We thank Dr D. Gordon for assistance in the identification of the bryozoan *G. mimicus*

and Dr B. Hayward for returning from ODP 181 with the sediment sample which has been made freely available to us. Figure 1 was drafted by L. Cotterall.

References

- Barnes, D.K.A. & Sanderson, W.G. (2000). Latitudinal patterns in the colonization of marine debris. In *Proceedings of the 11th International Bryozoology Association Conference, Smithsonian Tropical Research Institute, Panama*. (eds. Herrera-Cubilla, A. & Jackson, J.B.C.), pp154-160.
- Backhurst, M.K. & Cole, R.G. (2000). Subtidal benthic marine litter at Kawau Island, north-eastern New Zealand. *Journal of Environmental Management* 60: 227-237.
- Bingel, F., Avsar, D. & Unsal, M. (1987). A note on plastic materials in trawl catches in the north-eastern Mediterranean. *Meeresforschung* 31: 227-233.
- Coe, J.E. & Rogers, D.B. (eds.) (1997). *Marine Debris: sources, impacts, and solutions*. New York. Springer. 432p.
- Galgani, F., Burgeot, T., Bocquene, G., Vincent, F., Leaute J.P., Labastie, J., Forrest, A. & Guichet, R. (1995a). Distribution and abundance of debris on the continental shelf of the Bay of Biscay and in Seine Bay. *Marine Pollution Bulletin* 30: 58-62.
- Galgani, F., Jaunet, S., Campillo, A., Guenegen, X. & His, E. (1995b). Distribution and abundance of debris on the continental shelf of the north-western Mediterranean Sea. *Marine Pollution Bulletin* 30: 713-717.
- Galgani, F., Leaute, J.P., Moguedet, P., Souplets, A., Verin, Y., Carpentier, A., Gourageur, H., Latrouite, D., Andral, B., Cadiou, Y., Mahe, J.C., Poulard, J.C., & Nerisson, P. 2000. Litter on the sea floor along European coasts. *Marine Pollution Bulletin* 40: 516-527.
- Galgani, F., Souplet, A. & Cadiou, Y. (1996). Accumulation of debris on the deep sea floor off the French Mediterranean coast. *Marine Ecology Progress Series* 142: 225-234.
- Galil, B.S., Golik, A. & Turkay, M. (1995). Litter at the bottom of the sea: a sea bed survey in the eastern Mediterranean. *Marine Pollution Bulletin* 30: 22-24.
- Goldberg, E.D. (1997). Plasticizing the sea floor: an overview. *Environmental Technology*, 18: 195-202.
- Gordon, D.P. (1989). The Marine Fauna of New Zealand: Bryozoa: Gymnolaemata (Cheilostomida Ascophorina) from the western South Island continental shelf and slope. New Zealand Oceanographic Institute Memoir 97. 158p.
- Gordon, D.P. & d'Hondt, J.-L. (1997). Bryozoa: Lepraliomorpha and other Ascophorina, mainly from New Caledonian waters. *Memoirs du Museum National d'Histoire Naturelle* 176: 9-124.
- Gregory, M.R. (1978). Accumulation and distribution of virgin plastic granules on New Zealand beaches. *New Zealand Journal of Marine and Freshwater Research* 12: 399-414.
- Gregory, M.R. (1998). Pelagic plastics and marine invaders. *Aliens* 7: 6-7
- Gregory, M.R. (2004). Marine debris: hangers-on and hitch-hiking aliens. Derelict fishing gear and related marine debris: an educational outreach seminar among APEC partners, January 13-16, 2004 Honolulu, Hawaii, Seminar Proceedings, 40-44.
- Gregory, M.R. & Ryan, P.G. (1997). Pelagic plastics and other persistent synthetic debris: a review of Southern

- Hemisphere perspectives. In *Marine Debris: sources, impacts, and solution* (eds. Coe, J.M. & Rogers, D.B.), pp 49-66. New York, Springer..
- Harms, J. (1990). Marine plastic litter as an artificial hard bottom fouling ground. *Helgoländer Meeresuntersuchungen* 44: 503-506.
- Hess, N.A., Ribic, C.A. & Vining, I. (1999). Benthic marine debris, with an emphasis on fishery-related items, surrounding Kodiak Island, Alaska, 1994-1996. *Marine Pollution Bulletin* 38: 885-890.
- Holmström, A. (1975). Plastic films on the bottom of the Skagerack. *Nature* 255: 622-623.
- Kanehiro, H., Tokai, T. & Matsuda, K. (1996). The distribution of litter in fishing ground of Tokyo Bay. *Fisheries Engineering* 32: 211-217.
- Klauswitz, W. (1984). Kunststoffe an der Küste und im mer: ein ökologisches problem. *Natur und Museum* 114(6): 162-174.
- Lenihan, H.S., Oliver, J.S., Oakden, J.M. & Stephenson, M.D. (1990). Intense and localised benthic marine pollution around McMurdo Station, Antarctica. *Marine Pollution Bulletin* 21: 422-430.
- Moore, S.L. and Allen, M.J. (2000). Distribution of anthropogenic and natural debris on the mainland shelf of the Southern California Bight. *Marine Pollution Bulletin* 40: 83-88.
- Oshima, S. (2000). Towards a "visual sea". *Hydro International* 4(5): 73.
- Powlik, J.J. (1995). The effects of discarded plastic on the colonization of nearshore substrata. *Sarsia* 80: 229-236.
- Stefatos, A., Charalampakis, M., Papatheodorou, G. & Ferentinos, G. (1999). Marine debris on the seafloor of the Mediterranean Sea: examples from two enclosed gulfs in western Greece. *Marine Pollution Bulletin* 36: 389-393.
- Stevens, L.M. (1992). Marine plastic debris: fouling and degradation. Unpublished MSc Thesis, University of Auckland. 110p.
- Stevens, L.M., Gregory, M.R. & Foster, B.A. (1996). Fouling bryozoa on pelagic and moored plastics from northern New Zealand. In *Bryozoans in space and time. Proceedings of the 10th International Bryozoology Conference, Wellington, New Zealand.* (eds. Gordon, D.P., Smith, A.M. & Grant-Mackie, J.A.), pp 321-340. National Institute of Water & Atmosphere Research Ltd, Wellington.
- Thompson, R.C., Olsen, Y., Mitchell, R.P., Davis, A., Rowland, S.J., John, A.W.G., McGonigle, D. & Russell, A.E. (2004). Lost at sea: where is all the plastic. *Science* 304: 838.
- Uneputty, P. & Evans, S.M. (1997). The impact of plastic debris on the biota of tidal flats in Ambon Bay (eastern Indonesia). *Marine Environmental Research* 44: 233-242.
- Williams, A.T., Simmons, S.L. & Fricker, A. (1993). Off-shore sinks of marine litter: a new problem. *Marine Pollution Bulletin* 26: 404-405.
- Winston, J.E. (1982). Drift plastic - an expanding niche for a marine invertebrate? *Marine Pollution Bulletin* 13: 348-351.
- Winston, J.E., Gregory, M.R. & Stevens, L.M. (1997). Encrusters, epibionts, and other biota associated with pelagic plastics: a review of biogeographical, environmental, and conservation issues. In *Marine Debris: sources, impacts, and solutions* (eds. Coe, J.M. & Rogers, D.B.), pp81-97. New York, Springer..
- Ye, S. & Andrady, A.L. (1991). Fouling

of floating plastic debris under Biscayne Bay exposure conditions. *Marine Pollution Bulletin* 22: 608-613.